


AD-A148 783


MULTIPHOTON PHOTOCHEMICAL AND COLLISIONAL EFFECTS
DURING OXYGEN ATOM FLAM. (U) ARMY BALLISTIC RESEARCH
LAB ABERDEEN PROVING GROUND MD A W NIZIOLEK ET AL
OCT 84 RRL-TR-2604 SBI-AD-F300 512 F/G 7/5

1/1

UNCLASSIFIED

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

**B
R
L**

AD-A148 783

ADF300512

TECHNICAL REPORT BRL-TR-2604

**MULTIPHOTON PHOTOCHEMICAL AND
COLLISIONAL EFFECTS DURING OXYGEN
ATOM FLAME DETECTION**

**Andrzej W. Miziolek
Mark A. DeWilde**

October 1984

DTIC FILE COPY

**DTIC
ELECTE
NOV 29 1984
S D E**

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

**US ARMY BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MARYLAND**

84 11 27 055

**Destroy this report when it is no longer needed.
Do not return it to the originator.**

**Additional copies of this report may be obtained
from the National Technical Information Service,
U. S. Department of Commerce, Springfield, Virginia
22161.**

**The findings in this report are not to be construed as an official
Department of the Army position, unless so designated by other
authorized documents.**

**The use of trade names or manufacturers' names in this report
does not constitute indorsement of any commercial product.**

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
TECHNICAL REPORT BRL-TR-2604	AD-A148 783	
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
MULTIPHOTON PHOTOCHEMICAL AND COLLISIONAL EFFECTS DURING OXYGEN ATOM FLAME DETECTION		Final
6. PERFORMING ORG. REPORT NUMBER		
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
Andrzej W. Miziolek Mark A. DeKilde		
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBER
US Army Ballistic Research Laboratory ATTN: AMXBR - IBD Aberdeen Proving Ground, MD 21005-5066		1L161102AH43
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
US Army Ballistic Research Laboratory ATTN: AMXBR-OD-ST Aberdeen Proving Ground, MD 21005-5066		October 1984
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES
		23
		15. SECURITY CLASS (of this report)
		Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release; distribution is unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Photochemistry Multiphoton Fluorescence Energy Transfer Oxygen Detection Flame Atoms		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
meg A Nd:YAG pumped dye laser system was used to two-photon excite oxygen atoms at 225.6 nm in an atmospheric pressure $\text{CH}_4/\text{N}_2\text{O}/\text{N}_2$ flame. Subsequent emission at 844.7 nm from the directly populated state, as well as a stronger emission at 777.5 nm due to the $\text{O}(3p^3\text{P} \rightarrow 3p^5\text{P})$ collisional energy transfer process, was monitored. Two-photon resonant oxygen atom and hydrogen atom (656.3 nm) emissions were also observed in the absence of a flame. Closer examination revealed that the tightly focussed probe beam was producing		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. Abstract (Cont'd):

these atoms by promoting multiphoton photolysis of the oxidizer, as well as fuel molecules. Thus, this type of laser diagnostic probe is potentially quite intrusive depending on the combustion region that is probed, as well as the laser energies used.


UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

TABLE OF CONTENTS

	<u>Page</u>
I. INTRODUCTION.....	5
II. EXPERIMENTAL.....	6
III. RESULTS AND DISCUSSION.....	6
IV. CONCLUSIONS.....	13
ACKNOWLEDGEMENTS.....	14
REFERENCES.....	15
DISTRIBUTION LIST.....	17

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



I. INTRODUCTION

Propellant combustion is a process of considerable complexity in which high temperature chemical reactions play a major role. In order to understand the actual heat releasing chemical steps that are involved, an extensive program has been ongoing at BRL, as well as in other combustion research institutions, where much simpler laboratory model flames are studied in detail. Such studies involve the interplay of flame diagnostics (usually based on laser spectroscopy techniques) with flame modeling work, as well as allied kinetics efforts to supply necessary elementary reaction rate constants for such detailed chemical flame models. In the area of flame diagnostics, laser spectroscopy techniques such as absorption, Raman, laser induced fluorescence (LIF), coherent anti-Stokes Raman spectroscopy (CARS), etc., have been able to detect the stable major flame species as well as many reactive molecular intermediates. There is, however, an important class of combustion intermediates, i.e., the flame atoms, such as H, O, and N, which has historically eluded direct detection due to the lack of accessible energy levels for single photon excitation.

Very recently, though, multiphoton excitation techniques have been successful in detecting the hydrogen and oxygen atoms in a flame. So far, two distinct detection approaches have been demonstrated, one involving multiphoton emission (MPE)^{1,2} and the other multiphoton ionization (MPI),^{3,4,5} the latter also referred to as optogalvanic detection. In the case of the oxygen atoms, however, direct detection in a flame was first accomplished by spontaneous Raman scattering⁶ and coherent anti-Stokes Raman spectroscopy (CARS).⁷ Apparently, sensitivity limits and spectral interferences have prevented further development of these spectroscopies for atomic flame detection.

¹M. Alden, H. Edner, P. Grafstrom, and S. Svanberg, "Two-Photon Excitation of Atomic Oxygen in a Flame," *Opt. Comm.*, Vol. 42, p. 244, 1982.

²R.P. Lucht, J.T. Salmon, G.B. King, D.W. Sweeney, and N.M. Laurendeau, "Two-Photon-Excited Fluorescence Measurement of Hydrogen Atoms in Flames," *Opt. Lett.*, Vol. 8, p. 365, 1983.

³J.E.M. Goldsmith, "Resonant Multiphoton Optogalvanic Detection of Atomic Hydrogen in Flames," *Opt. Lett.*, Vol. 7, p. 437, 1982.

⁴J.E.M. Goldsmith, "Resonant Multiphoton Optogalvanic Detection of Atomic Oxygen in Flames," *J. Chem. Phys.*, Vol. 78, p. 1610, 1983.

⁵P.J.H. Tjossen and T.A. Cool, "Detection of Atomic Hydrogen in Flames by Resonance Four-Photon Ionization at 365 nm," *Chem. Phys. Lett.*, Vol. 100, p. 479, 1983.

⁶G.J. Dasch and J.H. Bechtel, "Spontaneous Raman Scattering by Ground-State Oxygen Atoms," *Opt. Lett.*, Vol. 6, p. 36, 1981.

⁷R.E. Teets and J.H. Bechtel, "Coherent Anti-Stokes Raman Spectra of Oxygen Atoms in Flames," *Opt. Lett.*, Vol. 6, p. 458, 1981.

Both the MPI and the MPE approaches suffer certain limitations: MPE always has quenching considerations as well as photometric inefficiencies, while MPI lacks selectivity in signal detection, and is thus quite susceptible to background ionization. In either approach, the nonlinear signals which usually depend on the number of photons, n , required for the given transition, are generally low due to the characteristically small multiphoton absorption cross sections. Thus, tight focusing and maximum laser energy extraction is typically desirable. We have made recent observations in our laboratory, however, which suggest that at sufficiently high laser energies, the flame diagnostic probe can become substantially intrusive by promoting multiphoton dissociation of the parent fuel and oxidizer molecules, in this case into hydrogen and oxygen atoms. Also, we have observed very efficient collisional energy transfer from the two-photon populated oxygen level to a neighboring level, as well as energy transfer from the excited oxygen to hydrogen atoms (see Figure 1). The two-photon excitation scheme for oxygen atoms using 225 nm photons, followed by detection of emission at 845 nm (see Figure 1), was initially demonstrated in a low pressure discharge⁸ and then applied successfully for the first time to a combustion environment in an experiment involving the C_2H_2/O_2 flame.¹

II. EXPERIMENTAL

Figure 2 shows the experimental schematic. A Nd:YAG laser is used to pump a dye laser in which R590 dye is circulated. The output is frequency doubled and mixed with the 1.06 micron Nd:YAG fundamental in a commercial wavelength extender to produce tunable radiation in the 225 nm range. This beam is then focussed with a 100 mm focal length lens into a curved knife-edge burner, whose edges are separated by 3 mm from each other, and in which a fuel rich $CH_4/N_2O/N_2$ flame (0.14:0.38:0.48) is stabilized. Laser pulse widths of 5 nsec and energies up to 1.5 mJoules/pulse at 10 pps were employed. The emitted radiation from the flame was collected by a lens system and passed through a broadband filter into a 0.3 m monochromator whose slits were relatively wide (ca. 2 nm) and finally detected by a photomultiplier tube (EMI 9659QA). The PMT output was fed into a 7912AD Tektronix transient digitizer, which was interfaced with a PDP 11/04 computer. Typically 50-300 laser shots were averaged per data point.

III. RESULTS AND DISCUSSION

Figure 3 shows an O atom emission profile through the curved knife-edge burner. A well-defined flame front is quite evident while the second peak is interpreted as resulting from entrained air further reacting with the fuel-rich post-flame gases. Figure 3 is based on monitoring the 777.5 nm line, which is actually a grouping of three closely spaced lines originating from the three 5P upper states to the common 5S_0 state. The emission intensity of this group as measured in the primary flame zone was about an order of

⁸ W.K. Biachol, B.E. Perry, and D.R. Crosley, "Two-Photon Laser-Induced Fluorescence in Oxygen and Nitrogen Atoms," *Chem. Phys. Lett.*, Vol. 82, p. 85, 1981.






Figure 2. Experimental Schematic for O-Atom MPE Experiments




Figure 3. O Atom Emission Profile at 777.5 nm in a Curved Knife-Edge Burner Supporting a Fuel-Rich $\text{CH}_4/\text{N}_2\text{O}/\text{N}_2$ Flame

magnitude greater than the emission at 844.7 nm. When photometric wavelength differences, such as grating and PMT quantum efficiencies are taken into account, the 777.5/844.7 emission ratio is still about 3. This indicates an efficient energy transfer process and has desirable analytical implications since now two different wavelengths are available for O atom detection. Due to the fact that low signal levels were usually a problem, most of the experiments were done monitoring the 777.5 nm line.

Figure 4 shows the O atom two-photon excitation spectra where the ground spin-orbit states are resolved. The observed ratio of 9.5:4.1:1 can be compared to a Boltzmann calculated ratio of 5.8:3.1:1 for an adiabatic flame temperature of 2290°K. This, in turn, can be compared to the ratio of 1.6:1.3:1, which was observed in optogalvanic experiments on the H₂/O₂/Ar flame.⁴ The reason for these discrepancies is not clear at this time, but recently there have been further theoretical efforts to substantiate the two-photon absorption cross sections which were found to be equal for each of the three fine-structure components involved in this transition.⁹ A rough estimate of 0.0001% for the detection limit is based on the emission intensity in the primary flame zone (see Figure 3) and a calculated O atom mole fraction of 0.005% (NASA/Lewis equilibrium program). This number should be improved by at least an order of magnitude by optimizing the optical collection efficiency, as well as using a red-sensitive PMT. This detection limit can be compared to 0.01% indicated in the previous MPE experiment,¹ parts-per-million sensitivity for the MPI experiment,⁴ as well as 0.01-0.1% for the two Raman experiments.^{6,7}

Further spectral investigations revealed another emission line centered at 656.3 nm, which was generally as intense as the 777.5 nm line and was resonant with O atom two-photon excitation. We have identified this emission as the H_α transition in the hydrogen atom Balmer series (see Figure 1). However, the mechanism for populating the n=3 level, which is 8861 cm⁻¹ higher in energy than the laser populated O atom level, was quite unclear for some time. Another even more surprising observation was the detection of the two O atom emission lines, as well as the 656.3 nm H-atom line in the absence of a flame when only room temperature premixed gases were flowing through the burner. In order to understand these findings, a determination of the order of nonlinearity, n, of these signals under different conditions was initiated and the results are given in Table 1. The uncertainties for the value of n are based on a number of different runs in which the Nd:YAG amplifier lamp energy was varied to yield energies between 0.2-1.5 mJoule/pulse. Clearly, this is not the best way to vary the probe laser energy since the output pulse time profile may change somewhat. Unfortunately, intensity attenuation attempts using polarizers available in our laboratory were unsuccessful since the coatings could not withstand the high peak intensities and short wavelengths used. Nevertheless, the results listed in Table 1 are consistent with a qualitative picture of multiphoton induced photolysis of the fuel and oxidizer parent molecules followed by a O atom two-photon resonant formation of a microplasma. In the preheat region the value of n=4 is close to what is found for the N₂O room temperature flow where n=5, suggesting that multiphoton

⁹D.R. Crosley and W.K. Bischel, "On Relative Fine-Structure Intensities in Two-Photon Excitation," *Phys. Rev.*, in press 1984.




Figure 4. O-Atom Emission at 777.5 nm Due to Spin-Orbit Resolved Ground States in the Primary Flame Zone

TABLE 1. ORDER OF NONLINEARITY (n) OF MULTIPHOTON INDUCED EMISSION SIGNALS^a

Atom (Emission λ)	Flame Region Probed	n
O (777.5 nm)	Preheat	4 ± 0.5
"	Primary	2 ± 0.5
"	Post-Flame	1.8 ± 0.5
"	Room Temp. Flow	5 ± 1
H (656.3 nm)	"	$10 +$

a - Where emission signal \propto (laser energy)ⁿ

photolysis of N₂O is contributing significantly to the total O atom emission signal. In the primary and post-flame regions, however, the value of n=2 is just what one would expect for a two-photon generated signal where the photolytic parent, N₂O, has already reacted away and the native O atom flame population is relatively high. The highest order of nonlinearity is found for the H atom emission which is expected since it takes some number of photons to photolyze CH₄, and more for N₂O, and finally two to excite the O atoms. The lack of uncertainty limits for the H-atom case attests to the difficulty (i.e., long averaging times and high degree of scatter) involved in making measurements on very high orders of nonlinearity.

A microplasma is not surprising since the absorption of the third photon ionizes the O atoms⁴ and the temperature in the focal volume may be much hotter than even the flame temperature, thus promoting collisional energy transfer from O to H atoms. An alternative to multiphoton photolysis might be plasma chemistry involving highly excited states or ion-molecule reactions, but these explanations seem less likely since all of the emissions are prompt, i.e., 5 nsec or less apart from each other. Again, it should be stressed that the results in Table 1 are qualitative in nature since signal levels, especially where n is high, were quite low, requiring laser energies over 1 mJoule/pulse, and they are based on the collisionally induced 777.5 nm line where the rates of collisional energy transfer may not be constant for the different experimental conditions. Clearly, this is a complex situation which deserves further study. Nevertheless, the potential for the multiphoton diagnostic probe to be intrusive certainly exists and thus appropriate caution should be exercised in these types of experiments. The implication of our results to previous multiphoton O atom flame studies cannot be readily assessed since those were done on flame systems other than the CH₄/N₂O/N₂ and thus photochemical effects (if any) are probably quite different. Finally, multiphoton induced photolysis should, in fact, not be too surprising since recent literature in chemical physics has many examples of such, even

involving typical flame molecules like H_2 ¹⁰ or C_2H_2 .¹¹ There is even a recent flame paper where laser probe intrusion (presumably photochemical) was noted.¹² On the other hand, as pointed out by those authors, there may be new opportunities to expand our knowledge of combustion by exploiting photochemistry.

IV. CONCLUSIONS

The following conclusions can be reached on the basis of our experiments.

1. Oxygen atoms can be detected in a $CH_4/N_2O/N_2$ flame by two-photon excitation at 225.6 nm and emission detection at 844.7 nm.

2. The collisional energy transfer between the directly populated 3P excited oxygen atom state and the neighboring 5P state is very efficient. This is quite surprising since the process involves the change of spin multiplicities. For a given experimental condition, the resulting emission at 777.5 nm can be much more intense than the one at 844.7 nm and thus be more useful for sensitive analysis of O-atoms.

3. Hydrogen atom emission at 656.3 nm is detected, which is resonant with O atom two-photon excitation. This may result from an endothermic collisional energy transfer process occurring in the microplasma formed in the focal volume of the laser whose temperature is hotter than the flame temperature. Alternative explanations, however, for the H-atom emission such as plasma chemistry/fast ion-molecule reactions, cannot be ruled out at this time.

4. Since both oxygen atom and the single hydrogen atom emissions were detected without a flame, the multiphoton photolysis of the atom precursors, N_2O and CH_4 , was indicated. This was substantiated by a study of the order of nonlinearity of the emission signal.

5. The above observation indicates that the laser multiphoton flame diagnostic probe is potentially quite intrusive, depending on the region of the flame being studied (i.e., the relative concentration of photochemical precursors), as well as the laser energies utilized (i.e., the higher the laser energy, the more prominent the higher order processes become). Thus, appropriate caution should be exercised in these kinds of experiments.

¹⁰S.T. Pratt, P.M. Dehmer, and J.L. Dehmer, "Resonant Multiphoton Ionization of H_2 via the $B^1\Sigma_u^+$, $v=7$, $J=2$, and 4 Levels with Photoelectron Energy Analysis," *J. Chem. Phys.*, Vol. 78, p. 4315, 1983.

¹¹B.B. Craig, W.L. Faust, L.S. Goldberg, and R.G. Weiss, "UV Short-Pulse Fragmentation of Isotopically Labeled Acetylene: Studies of Emission with Subnanosecond Resolution," *J. Chem. Phys.*, Vol. 76, p. 5014, 1982.

¹²M. Alden, H. Edner, and S. Svanberg, "Simultaneous Spatially Resolved Monitoring of C_2 and OH in a C_2H_2/O_2 Flame Using a Diode Array Detector," *Appl. Phys. B*, Vol. 29, p. 93, 1982.

ACKNOWLEDGEMENTS

The authors would like to express their gratitude to the BRL Director, Dr. R.J. Eichelberger, for supporting this research under the ILIR Program. Also, numerous discussions with Professors Paul J. Dagdigian, Johns Hopkins University, and Millard H. Alexander, University of Maryland, College Park, were quite stimulating and helpful.

REFERENCES

1. M. Alden, H. Edner, P. Grafstrom, and S. Svanberg, "Two-Photon Excitation of Atomic Oxygen in a Flame," Opt. Comm., Vol. 42, p. 244, 1982.
2. R.P. Lucht, J.T. Salmon, G.B. King, D.W. Sweeney, and N.M. Laurendeau, "Two-Photon-Excited Fluorescence Measurement of Hydrogen Atoms in Flames," Opt. Lett., Vol. 8, p. 365, 1983.
3. J.E.M. Goldsmith, "Resonant Multiphoton Optogalvanic Detection of Atomic Hydrogen in Flames," Opt. Lett., Vol 7, p. 437, 1982.
4. J.E.M. Goldsmith, "Resonant Multiphoton Optogalvanic Detection of Atomic Oxygen in Flames," J. Chem. Phys., Vol. 78, p. 1610, 1983.
5. P.J.H. Tjossem and T.A. Cool, "Detection of Atomic Hydrogen in Flames by Resonance Four-Photon Ionization at 365 nm," Chem. Phys. Lett., Vol. 100, p. 479, 1983.
6. C.J. Dasch and J.H. Bechtel, "Spontaneous Raman Scattering by Ground-State Oxygen Atoms," Opt. Lett., Vol. 6, p. 36, 1981.
7. R.E. Teets and J.H. Bechtel, "Coherent Anti-Stokes Raman Spectra of Oxygen Atoms in Flames," Opt. Lett., Vol. 6, p. 458, 1981.
8. W.K. Bischel, B.E. Perry, and D.R. Crosley, "Two-Photon Laser-Induced Fluorescence in Oxygen and Nitrogen Atoms," Chem. Phys. Lett., Vol. 82, p. 85, 1981.
9. D.R. Crosley and W.K. Bischel, "On Relative Fine-Structure Intensities in Two-Photon Excitation," Phys. Rev., in press 1984.
10. S.T. Pratt, P.M. Dehmer, and J.L. Dehmer, "Resonant Multiphoton Ionization of H_2 via the $B^1 \Sigma_u^+$, $v=7$, $J=2$, and 4 Levels with Photoelectron Energy Analysis," J. Chem. Phys., Vol. 78, p. 4315, 1983.
11. B.B. Craig, W.L. Faust, L.S. Goldberg, and R.G. Weiss, "UV Short-Pulse Fragmentation of Isotopically Labeled Acetylene: Studies of Emission with Subnanosecond Resolution," J. Chem. Phys., Vol. 76, p. 5014, 1982.
12. M. Alden, H. Edner, and S. Svanberg, "Simultaneous Spatially Resolved Monitoring of C_2 and OH in a C_2H_2/O_2 Flame Using a Diode Array Detector," Appl. Phys. B, Vol. 29, p. 93, 1982.

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
12	Administrator Defense Technical Info Center ATTN: DTIC-DDA Cameron Station Alexandria, VA 22314	1	Director USA Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035
1	HQ DA DAMA-ART-M Washington, DC 20310	4	Commander US Army Research Office ATTN: R. Hhirardelli D. Mann R. Singleton D. Squire P. O. Box 12211 Research Triangle Park, NC 27709
1	Commander US Army Materiel Command ATTN: AMCDRA-ST 5001 Eisenhower Avenue Alexandria, VA 22333	1	Commander USA Communications Research and Development Command ATTN: AMSEL -ATDD Fort Monmouth, NJ 07703
1	Commander Armament R&D Center USA AMCCOM ATTN: SMCAR-TDC Dover, NJ 07801	1	Commander USA Electronics Research and Development Command Technical Support Activity ATTN: AMDSD -L Fort Monmouth, NJ 07703
1	Commander Armament R&D Center USA AMCCOM ATTN: SMCAR-TSS Dover, NJ 07801	2	Commander USA AMCCOM ATTN: SMCAR -LCA-G , D.S. Downs J.A. Lannon Dover, NJ 07801
1	Commander USA AMCCOM ATTN: AMSMC -LEP-L Rock Island, IL 61299	1	Commander USA AMCCOM ATTN: SMCAR -LC , L. Harris Dover, NJ 07801
1	Director Benet Weapons Laboratory Armament R&D Center USA AMCCOM ATTN: SMCAR-LCB-TL Watervliet, NY 12189	1	Commander USA AMCCOM ATTN: SMCAR-SCM-T, L. Stiefel Dover, NJ 07801
1	Commander USA Aviation Research and Development Command ATTN: AMSAV-E 4300 Goodfellow Blvd. St. Louis, MO 63120		

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
1	Commander USA Missile Command ATTN: AMSMI-R Redstone Arsenal, AL 35898	1	Commander Naval Air Systems Command ATTN: J. Ramnarace, AIR-54111C Washington, DC 20360
1	Commander USA Missile Command ATTN: AMSMI-YDL Redstone Arsenal, AL 35898	3	Commander Naval Ordnance Station ATTN: C. Irish S. Mitchell P.L. Stang, Code 515 Indian Head, MD 20640
2	Commander USA Missile Command ATTN: AMSMI-RK, D.J. Ifshin Redstone Arsenal, AL 35898	1	Commander Naval Surface Weapons Center ATTN: J.L. East, Jr., G-20 Dahlgren, VA 22448
1	Commander USA Tank Automotive Command ATTN: AMSTA-TSL Warren, MI 48090	2	Commander Naval Surface Weapons Center ATTN: R. Bernecker, R-13 G.B. Wilmot, R-16 Silver Spring, MD 20910
1	Director USA TRADOC Systems Analysis Activity ATTN: ATAA-SL WSMR, NM 88002	4	Commander Naval Weapons Center ATTN: R.L. Derr, Code 389 China Lake, CA 93555
1	Commandant US Army Infantry School ATTN: ATSH-CD-CSO-OR Fort Benning, GA 31905	2	Commander Naval Weapons Center ATTN: Code 3891, T. Boggs K.J. Graham China Lake, CA 93555
1	Commander US Army Development and Employment Agency ATTN: MODE-TED-SAB Fort Lewis, WA 98433	5	Commander Naval Research Laboratory ATTN: L. Harvey J. McDonald E. Oran J. Shnur R.J. Doyle, Code 6110 Washington, DC 20375
1	Office of Naval Research Department of the Navy ATTN: R.S. Miller, Code 432 800 N. Quincy Street Arlington, VA 22217	1	Commander Naval Underwater Systems Center Weapons Dept. ATTN: R.S. Lazar/Code 36301 Newport, RI 02840
1	Navy Strategic Systems Project Office ATTN: R.D. Kinert, SP 2731 Washington, DC 20376		

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
1	Superintendent Naval Postgraduate School Dept. of Aeronautics ATTN: D.W. Netzer Monterey, CA 93940	2	Atlantic Research Corp. ATTN: M.K. King 5390 Cherokee Avenue Alexandria, VA 22314
6	AFRPL (DRSC) ATTN: R. Geisler D. George B. Goshgarian J. Levine W. Roe D. Weaver Edwards AFB, CA 93523	1	Atlantic Research Corp. ATTN: R.H.W. Waesche 7511 Wellington Road Gainesville, VA 22065
1	ADTC/AFATL/DLDL ATTN: O.K. Heiney Eglin AFB, FL 32542	1	AVCO Everett Rsch. Lab. Div. ATTN: D. Stickler 2385 Revere Beach Parkway Everett, MA 02149
1	AFOSR ATTN: L.H. Caveny Bolling Air Force Base Washington, DC 20332	1	Battelle Memorial Institute Tactical Technology Center ATTN: J. Huggins 505 King Avenue Columbus, OH 43201
1	AFWL/SUL Kirtland AFB, NM 87117	2	Exxon Research & Eng. Co. ATTN: A. Dean M. Chou P.O. Box 45 Linden, NJ 07036
1	Director NASA Langley Research Center ATTN: G.B. Northam/MS 168 Hampton, VA 23365	1	Ford Aerospace and Communications Corp. DIVAD Division Div. Hq., Irvine ATTN: D. Williams Main Street & Ford Road Newport Beach, CA 92663
4	National Bureau of Standards ATTN: J. Hastie M. Jacox T. Kashiwagi H. Semerjian US Department of Commerce Washington, DC 20234	1	General Electric Armament & Electrical Systems ATTN: M.J. Bulman Lakeside Avenue Burlington, VT 05401
1	Aerojet Solid Propulsion Co. ATTN: P. Micheli Sacramento, CA 95813	1	General Electric Company ATTN: M. Lapp 2352 Jade Lane Schenectady, NY 12309
1	Applied Combustion Technology, Inc. ATTN: A.M. Varney P.O. Box 17885 Orlando, FL 32860		

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
1	General Motors Rsch Labs Physics Department ATTN: J.H. Bechtel Warren, MI 48090	1	Los Alamos National Lab ATTN: B. Nichols T7, MS-B284 P.O. Box 1663 Los Alamos, NM 87544
3	Hercules, Inc. Allegheny Ballistics Lab. ATTN: R.R. Miller P.O. Box 210 Cumberland, MD 21501	1	Olin Corporation Smokeless Powder Operations ATTN: R.L. Cook P.O. Box 222 St. Marks, FL 32355
3	Hercules, Inc. Bacchus Works ATTN: K.P. McCarty P.O. Box 98 Magna, UT 84044	1	Paul Gough Associates, Inc. ATTN: P.S. Gough 1048 South Street Portsmouth, NH 03801
1	Hercules, Inc. AFATL/DLDL ATTN: R.L. Simmons Eglin AFB, FL 32542	2	Princeton Combustion Research Laboratories, Inc. ATTN: M. Summerfield N.A. Messina 475 US Highway One Monmouth Junction, NJ 08852
1	Honeywell, Inc. Defense Systems Division ATTN: D.E. Broden/ MS MN50-2000 600 2nd Street NE Hopkins, MN 55343	1	Hughes Aircraft Company ATTN: T.E. Ward 8433 Fallbrook Avenue Canoga Park, CA 91303
1	IBM Corporation ATTN: A.C. Tam Research Division 5600 Cottle Road San Jose, CA 95193	1	Rockwell International Corp. Rocketdyne Division ATTN: J.E. Flanagan/HB02 6633 Canoga Avenue Canoga Park, CA 91304
1	Director Lawrence Livermore Laboratory P.O. Box 808 ATTN: C. Westbrook Livermore, CA 94550	2	Sandia National Laboratories Combustion Sciences Dept. ATTN: R. Cattolica D. Stephenson P. O. Box 969 Livermore, CA 94554
1	Lockheed Missiles & Space Co. ATTN: George Lo 3251 Hanover Street Dept. 52-35/B204/2 Palo Alto, CA 94304	1	Sandia National Laboratories ATTN: M. Smooke Division 8353, P. O. BOX 969 Livermore, CA 94550
		1	Science Applications, Inc. ATTN: R.B. Edelman 23146 Cumorah Crest Woodland Hills, CA 91364

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
1	Science Applications, Inc. ATTN: H.S. Pergament 1100 State Road, Bldg. N Princeton, NJ 08540	2	United Technologies Corp. ATTN: R.S. Brown R.O. McLaren P.O. Box 358 Sunnyvale, CA 94086
1	United Technologies ATTN: A.C. Eckbreth East Hartford, CT 06108	1	Universal Propulsion Company ATTN: H.J. McSpadden Black Canyon Stage 1 Box 1140 Phoenix, AZ 85029
4	SRI International ATTN: S. Barker D. Crosley D. Golden Tech Lib 333 Ravenswood Avenue Menlo Park, CA 94025	1	Veritay Technology, Inc. ATTN: E.B. Fisher P.O. Box 22 Bowmansville, NY 14026
1	Stevens Institute of Tech. Davidson Laboratory ATTN: R. McAlevy, III Hoboken, NJ 07030	1	Brigham Young University Dept. of Chemical Engineering ATTN: M.W. Beckstead Provo, UT 84601
1	Teledyne McCormack-Selph ATTN: C. Leveritt 3601 Union Road Hollister, CA 95023	1	Director Jet Propulsion Laboratory ATTN: MS 125/159 4800 Oak Grove Drive Pasadena, CA 91109
1	Thiokol Corporation Elkton Division ATTN: W.N. Brundige P.O. Box 241 Elkton, MD 21921	1	California Institute of Technology ATTN: F.E.C. Culick/ MC 301-46 204 Karman Lab. Pasadena, CA 91125
3	Thiokol Corporation Huntsville Division ATTN: D.A. Flanagan Huntsville, AL 35807	1	University of California, Berkeley Mechanical Engineering Dept. ATTN: J. Daily Berkeley, CA 94720
3	Thiokol Corporation Wasatch Division ATTN: J.A. Peterson P.O. Box 524 Brigham City, UT 84302	1	University of California Los Alamos National Lab. ATTN: T.D. Butler P.O. Box 1663, Mail Stop B216 Los Alamos, NM 87544

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
2	University of California, Santa Barbara Quantum Institute ATTN: K. Schofield M. Steinberg Santa Barbara, CA 93106	1	University of Illinois Dept. of Mech. Eng. ATTN: H. Krier 144MEB, 1206 W. Green St. Urbana, IL 61801
1	University of Southern California Dept. of Chemistry ATTN: S. Benson Los Angeles, CA 90007	1	Johns Hopkins University/APL Chemical Propulsion Information Agency ATTN: T.W. Christian Johns Hopkins Road Laurel, MD 20707
1	Case Western Reserve Univ. Div. of Aerospace Sciences ATTN: J. Tien Cleveland, OH 44135	1	University of Minnesota Dept. of Mechanical Engineering ATTN: E. Fletcher Minneapolis, MN 55455
1	Cornell University Department of Chemistry ATTN: E. Grant Baker Laboratory Ithaca, NY 14850	4	Pennsylvania State University Applied Research Laboratory ATTN: G.M. Faeth K.K. Kuo H. Palmer M. Micci University Park, PA 16802
1	Univ. of Dayton Rsch Inst. ATTN: D. Campbell AFRPL/PAP Stop 24 Edwards AFB, CA 93523	1	Polytechnic Institute of NY ATTN: S. Lederman Route 110 Farmingdale, NY 11735
1	University of Florida Dept. of Chemistry ATTN: J. Winefordner Gainesville, FL 32601	2	Princeton University Forrestal Campus Library ATTN: K. Brezinsky I. Glassman P.O. Box 710 Princeton, NJ 08540
3	Georgia Institute of Technology School of Aerospace Engineering ATTN: E. Price Atlanta, GA 30332	1	Princeton University MAE Dept. ATTN: F.A. Williams Princeton, NJ 08544
2	Georgia Institute of Technology School of Aerospace Engineering ATTN: W.C. Strahle B.T. Zinn Atlanta, GA 30332		

DISTRIBUTION LIST

<u>No. Of</u> <u>Copies</u>	<u>Organization</u>	<u>No. Of</u> <u>Copies</u>	<u>Organization</u>
2	Purdue University School of Aeronautics and Astronautics ATTN: R. Glick J.R. Osborn Grissom Hall West Lafayette, IN 47906		<u>Aberdeen Proving Ground</u> Dir, USAMSAA ATTN: AMXSY -D AMXSY -MP, H. Cohen Cdr, USATECOM ATTN: AMSTE -TO-F Cdr, CRDC, AMCCOM ATTN: SMCCR-RSP-A SMCCR-MU SMCCR-SPS-IL
3	Purdue University School of Mechanical Engineering ATTN: N.M. Laurendeau S.N.B. Murthy D. Sweeney TSPC Chaffee Hall West Lafayette, IN 47906		
1	Rensselaer Polytechnic Inst. Dept. of Chemical Engineering ATTN: A. Fontijn Troy, NY 12181		
2	Southwest Research Institute ATTN: R.E. White A.B. Wenzel 8500 Culebra Road San Antonio, TX 78228		
1	Stanford University Dept. of Mechanical Engineering ATTN: R. Hanson Stanford, CA 94305		
2	University of Texas Dept. of Chemistry ATTN: W. Gardiner H. Schaefer Austin, TX 78712		
1	University of Utah Dept. of Chemical Engineering ATTN: G. Flandro Salt Lake City, UT 84112		
1	Virginia Polytechnic Institute and State University ATTN: J.A. Schetz Blacksburg, VA 24061		

USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. BRL Report Number _____ Date of Report _____
2. Date Report Received _____
3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) _____

4. How specifically, is the report being used? (Information source, design data, procedure, source of ideas, etc.) _____

5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided or efficiencies achieved, etc? If so, please elaborate. _____

6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) _____

CURRENT ADDRESS	_____
	Name

	Organization

	Address

	City, State, Zip

7. If indicating a Change of Address or Address Correction, please provide the New or Correct Address in Block 6 above and the Old or Incorrect address below.

OLD ADDRESS	_____
	Name

	Organization

	Address

	City, State, Zip

(Remove this sheet along the perforation, fold as indicated, staple or tape closed, and mail.)

----- FOLD HERE -----

Director
US Army Ballistic Research Laboratory
ATTN: AMXBR-OD-ST
Aberdeen Proving Ground, MD 21005-5066




NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

BUSINESS REPLY MAIL
FIRST CLASS PERMIT NO 12062 WASHINGTON, DC
POSTAGE WILL BE PAID BY DEPARTMENT OF THE ARMY

Director
US Army Ballistic Research Laboratory
ATTN: AMXBR-OD-ST
Aberdeen Proving Ground, MD 21005-9989



----- FOLD HERE -----

END

FILMED

1-85

DTIC